



## ALL Family Insect Repellent Spray 20 %

Saltidin® (known as Picaridin in the US)

Issued in February 2011

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## Summary

	Page
<b>1. Preliminary Remarks</b>	3
<b>2. Physical and Chemical Properties</b>	4
<b>3. Formulation</b>	6
3.1 Formulation	6
3.2 Preparation	6
3.3 Physical data	7
3.4 Packaging material	7
3.5 Stability	7
<b>4. Toxicological Data</b>	8
<b>5. Efficacy Studies</b>	9
5.1 Effectiveness against Mosquitoes	9
5.2 Effectiveness against Ticks	14
5.3 Effectiveness against Flies	16
5.4 Effectiveness against Sandflies	18
5.5 Effectiveness against Biting Midges	19
5.6 Effectiveness against Fleas	19
5.7 Effectiveness against Ants	20
5.8 Effectiveness against Cockroaches	21
<b>6. References</b>	22

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## 1. Preliminary Remarks

The use of repellents by application to skin and clothing is an important means of human protection against blood-sucking insects and ticks as well as arthropod-borne disease transmission.

Saltidin® belongs to a new generation of synthetic repellents developed as an alternative to DEET.

Developed by Bayer AG using molecular modelling techniques, Saltidin® showed the best performance among more than 800 substances against a variety of arthropods and the best characteristics as far as safety and compatibility with skin and plastic materials are concerned.

Presently, this molecule is owned by Saltigo GmbH (LANXESS Group) and is marketed by LANXESS Corporation (previously a Division of Bayer Corporation) in the USA.

Saltidin® is a safe and reliable insect repellent and stands for effective protection against various diseases transmitted by:

- ✓ Mosquitoes
- ✓ Ticks
- ✓ Flies
- ✓ Horseflies
- ✓ Sandflies
- ✓ Biting midges
- ✓ Fleas
- ✓ Ants
- ✓ Cockroaches

Due to its broad spectrum of efficacy, its favourable toxicological and cosmetic properties, Saltidin® can be safely used by consumers.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.




## 2. Physical and Chemical Properties

Chemical Name: 1-PIPERIDINECARBOXYLIC ACID, 2-(2-HYDROXY-ETHYL), 1-METHYLPROPYLESTER

Commercial Name: Saltidin®

Other Names: Icaridin, KBR 3023, Bayrepel™, Picaridin, (US EPA Registration)

Structure:



Empirical Formula: C<sub>12</sub>H<sub>23</sub>NO<sub>3</sub>

ELINCS Number 423-210-8

CAS Number 119515-38-7

Molecular weight: 229.3 g/mol

Melting point: < -170 °C

Boiling point: 272 °C at 1013 hPa

Viscosity: 30.7 sec. flow time accord. To DIN 53211, Dynamic: 129 mPa.s at 20 °C

Vapour pressure: 3.4 x 10<sup>-4</sup> hPa at 20 °C  
5.9 x 10<sup>-4</sup> hPa at 25 °C  
7.1 x 10<sup>-3</sup> hPa at 50 °C

Flash point: 142 °C

Density: 1.036 g/m<sup>3</sup> at 20 °C

Solubility: In water: 8.6 g/L at 20 °C  
In propane-2-ol: > 250 g/L at 20 °C  
In acetone: > 250 g/L at 20 °C  
In n-heptane: > 250 g/L at 20 °C  
In xylene: > 250 g/L at 20 °C

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



Appearance:	Clear liquid, colourless to brownish
Odour:	Slightly odorous
Storage conditions:	Store at room temperature

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



### 3. Formulation

#### ALL Family Insect Repellent Spray 20 %

##### 3.1 Formulation:

No	Ingredient	CAS no	Content %
1	Saltidin®	119515-38-7	20,6
2	Polyethylene Glycol 400	25322-68-3	20,0
3	Ethanol denatured	64-17-5	29,0
4	Floral D 12889 G	Not applicable	1,0
5	Water deionised		29,4

##### 3.2 Preparation:

One vessel manufacturing (closed vessel).

Mix all ingredients at room temperature and stir for 15 min. The sequence of ingredients is without relevance. Please note: Ethanol/water – light exothermic reaction.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



### 3.3 Physical data:

pH-value	7.0 - 9.0
Refractive Index	1.400 - 1.406
Oudor	Perfume conform
Appearance	Clear, light yellow liquid
Density	0.973 - 0.983
Content Saltidin® [%]	19 - 21

### 3.4 Packaging material:

	Volume	Material	Manufacturer
Bottle with pump valve	100ml	HDPE	Hermann Koch GmbH
Lotion bottle	100ml	HDPE	Hermann Koch GmbH

### 3.5 Stability

Long term GLP-stability for 3 years at 20 °C

Long term GLP-stability for 1 year at 25 °C/60 % rel. humidity

GLP - stability for 8 weeks 40 °C

Stability for 1 week at 0 °C

Stability for 2 weeks at 54 °C

The above formulation is intended solely as a guide for our business partners and others interested in our product. As the conditions of use and application of the suggested formulation are beyond our control, it is imperative that it be tested to determine, to your satisfaction, whether it is suitable for your intended use(s) and application(s). This application-specific analysis at least must include testing to determine suitability from a technical, as well as health, safety an environmental standpoints. Further, although the ingredients, quantities thereof and properties of compounds or finished goods mentioned herein reflect our recommendation at the time of publication, this guide may not be subject to continuous review and/or updating, and you agree that use is undertaken at your sole risk. All information is given without warranty or guarantee, and it is expressly understood and agreed that you assume, and hereby expressly release us from all liability, in fort contract or otherwise, incurred in connection with the use of this guide.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



#### 4. Toxicological Data

The toxicological properties of the proposed formulation have been assessed on the basis of available data, in accordance with the above intended usages on behalf of Saltigo GmbH (Germany). The studies are described in the report: "ICARIDIN. Tier II Summary Reports for Acute Studies All-Family Insect Repellent Spray (EP)", October 27, 2007.

<b>Extrapolated Acute Toxicity and Precautionary Signal Words for All-Family Insect Repellent Spray (20 % Spray-Formulation)</b>			
<b>Study type</b>	<b>Species</b>	<b>Results</b>	<b>EU Classification</b>
<b>Acute Oral</b>	Rat LD <sub>50</sub>	Male and Females = > 5050 mg/kg	Not classified
<b>Acute Dermal</b>	Rat LD <sub>50</sub>	Male and Females = > 5020 mg/kg	Not classified
<b>Acute Inhalation</b>	Rat LC <sub>50</sub>	> 30.2 mg/L (4- hours)	Not classified
<b>Eye Irritation</b>	Rabbit	No Eye Irritation	Not classified
<b>Skin Irritation</b>	Rabbit	No Skin Irritation	Not classified
<b>Dermal Sensitization</b>	Guinea Pig	Non Sensitizing	Not classified
<b>Skin Irritation</b>	Human	No Skin Irritation	Harmless

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## 5. Efficacy Studies

The efficacy of the proposed formulation has been assessed on the basis of available data, in accordance with the above intended usages on behalf of Saltigo GmbH (Germany). Reported here below are efficacy data against mosquitoes, ticks, flies, sandflies, biting midges and fleas obtained with formulations containing 10 - 20 % of Saltidin®. The studies are summarized in the report: "Summary Report of Efficacy Data Saltidin®", February 2011.

### 5.1 Effectiveness against Mosquitoes

Saltidin® has been shown to be a safe and effective repellent against various species of mosquitoes (e. g., *Aedes aegypti*, *Aedes albopictus*, *Culex quinquefasciatus*, and *Anopheles dirus*). Two types of protocols were used for testing the efficacy of Saltidin® on Mosquitoes: cage tests on human volunteers and field tests on human volunteers.

Cage tests on human volunteers were conducted during the development of the product to estimate the efficacy of the product and to compare different formulations of the product under standardized conditions.

The tests by Yap et al. (1997) were conducted in screened cages (1 x 1 x 1 m) with circular openings (15 cm in diameter) fitted with cloth sleeves. Eight volunteers (6 male and 2 females) were involved in the experiment. The application rate was 150 µL/90 cm<sup>2</sup>. The applied product was left to dry before covering it with a rubber sleeve with an opening window of 25 cm<sup>2</sup> as a surface for landing/biting activities of mosquitoes. Both hands of volunteers were covered with thick rubber gloves up to the wrists to prevent unwanted bites.

In the study by Luepkes (2005), the test protocol was similar to the Yap protocol: the cages measured 90 x 30 x 40 cm and the application rate was 150 µg/90 cm<sup>2</sup>.

The results showed that the 20 % Saltidin® formulation provided 4 - 8 hours protection and that the efficacy was similar or even superior to that of DEET (Table 1).

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



In the study by Zhao (2008) and Chen et al. (2008), the test protocol was similar to that of Yap. Results provided more than 7 and 10 hours protection against *Aedes albopictus* (Tiger Mosquito).

Table 1: Efficacy (Protection time = PT) of 20 % Saltidin® and DEET formulation against mosquitoes in cage tests.

Test Species	Test Location (Time of test)	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Aedes aegypti</i>	University Sains, Malaysia	20.0	4 - 6	20.0	4.0	Yap et al., 1997 <sup>[1]</sup>
<i>Aedes aegypti</i>	BioGenius GmbH, Monheim, Germany	20.0	6.8	Not determined		Luepkes, 2005 <sup>[2]</sup>
<i>Anopheles dirus</i>	University Sains, Malaysia	20.0	8.0	20.0	6.0	Yap et al., 1997 <sup>[1]</sup>
<i>Aedes albopictus</i>	Military Medical Research Institute, China	20.0	> 7.0	Not determined		Zhao, 2008 <sup>[3]*</sup>
<i>Aedes albopictus</i>	Nanjing Military District, Military Medical Research Institute, China	20.0	> 10	Not determined		Chen et al., 2008 <sup>[4]*</sup>

\* tests carried out on All Family Insect Repellent Spray 20 %

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



In the study by Prof. Dr. Mehlhorn (2010), the test protocol was similar to the Yap protocol: the cages measured 90 x 30 x 40 cm and the application rate was 150 µg/90 cm<sup>2</sup>.

The results showed that the 20 % Saltidin® formulation provided 4 - 6 hours protection against *Aedes aegypti* and that the efficacy was much better to that of IR3535 (Table 1a).

Table 1a: Efficacy (Protection time = PT) of 20 % Saltidin® and IR3535 formulation against mosquitoes in cage tests.

Test Species	Test Location (Time of test)	Saltidin®		IR3535		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Aedes aegypti</i>	Alpha-Biocare GmbH, Germany	20.0	4 - 6	20.0	1 - 2	Prof. Dr. Mehlhorn, 2010 <sup>[22]</sup>

Field tests were designed to determine the efficacy of products under real life conditions. It is common knowledge that cage tests are more conservative and that often the protection time for the cage tests is lower than in field tests. Therefore, field studies are important in order to establish the efficacy of the product. For this reason, this report will focus on field studies and on the more relevant data obtained from the cage tests using animals and human volunteers.

In the study by Yap et al. (1998), the trials against *Aedes albopictus* were conducted outdoor in a plantation on Penang Island during the day in order to coincide with the day-time biting activities. The trials against *Culex quinquefasciatus* were conducted in the living rooms of residential houses during the night to coincide with peak biting activities. The dosages tested of the repellent formulation were 1ml on the leg and 0.5 ml on the hand.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



In the studies by Frances et al. (2004), 4 male volunteers participated in the test. Each wore a long sleeved shirt buttoned at the wrist, long trousers and running shoes without socks. A mesh jacket was worn over the head and arms, and the legs of the trousers were rolled up to the knee to expose only the lower leg and ankles to biting mosquitoes. The repellents were spread evenly by the volunteers over each leg from the base of the knee to the ankle. The amount of repellent applied was determined by weighing the container before and after application. The mean of application rate was 0.91 - 1.67 mg/cm<sup>2</sup> for studies conducted in 2002 and 1.33 - 1.74 mg/cm<sup>2</sup> for studies in 2004.

In the study by Carroll-Loye Biological Research, carried out in Central Valley, California, 5 male volunteers and 5 female volunteers participated in the test. The application rates used in the field efficacy study were 0.97 µl/cm<sup>2</sup> on arms and 0.83 µl/cm<sup>2</sup> on legs and they exposed treated limbs to mosquitoes.

In the study against the mosquito species *Black fly* carried out by Carroll-Loye Biological Research 2010 in Southeastern California, 5 male volunteers and 5 female volunteers participated in the test. Each volunteer exposed one treated arm to wild population of *Black fly*.

The spray dosage rate of  $0.93 \pm 0.39$  mg/cm<sup>2</sup> on arms is the mean from the dosimetry analysis reported in related study. (Table 2 )

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



Table 2: Efficacy (Protection time = PT) of 20 % Saltidin® and DEET formulation against mosquitoes in field tests.

Test Species	Test Location (Time of test)	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Aedes albopictus</i>	Residential- Mainland Peninsular, Malaysia (Day)	20.0	8.0	20.0	6.0	Yap et al., 1998 <sup>[5]</sup>
<i>Culex quinquefasciatus</i>	Residential- Mainland Peninsular, Malaysia (Night)	20.0	8.0	20.0	8.0	Yap et al., 1998 <sup>[5]</sup>
<i>Mosquito spp</i>	Rain forest, Northern Queensland, Australia (Day, Night)	19.2	9.0	20.0	7.0	Frances et al., 2002 <sup>[6]</sup>
<i>Mosquito spp</i>	Northern Territory, Australia	19.2	8.0	20.0	8.0	Frances et al., 2004 <sup>[7]</sup>
<i>Mosquito spp</i>	Central Valley, California	20	11.6*	Not determined		Caroll-Loye, 2008 <sup>[8]</sup> *
<i>Black fly</i>	Central Valley, California	20	9.9*	Not determined		Caroll-Loye, 2010 <sup>[18]</sup> *

\* tests carried out on All Family Insect Repellent Spray 20 %

From the results of all the laboratory and field tests carried out by various researchers and in various parts of the world, it may be concluded that the formulation containing 20 % Saltidin® provides 8 - 11.6 hours of protection (Table 2). It can also be concluded that the performance of Saltidin® often proves to be superior to DEET in comparative studies.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## 5.2 Effectiveness against Ticks

Saltidin® has been shown to be a safe and effective repellent against various species of ticks (e. g., *Ixodes ricinus* and *Ixodes scapularis*).

In studies conducted in Austria by Sixl (1993), the results of cage tests and field tests on human volunteers showed that 20 % Saltidin® formulation provided at least four hours of protection (test duration) against *Ixodes ricinus*. A similar protection time was achieved using DEET.

In the field test, 20 people (9 men and 11 women) were involved in the experiment. The selected volunteers were particularly attractive to ticks. 10 - 14 ml of test formulations were applied to arms, legs and exposed areas of the body. The test duration was four hours (Table 3).

Table 3: Efficacy (Protection time = PT) of 20 % Saltidin® and DEET formulation against ticks in field tests.

Test Species	Test Location	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Ixodes ricinus</i>	Kaiser Wald near Graz, Austria	20.0	4.0**	20.0	4.0**	Sixl, 1993 <sup>[9]</sup>

\*\* Duration of the test

In the cage tests on human volunteers conducted in Austria by Sixl (1993), volunteers were treated with 20 % Saltidin® or 20 % DEET with protection extenders PEG400 and were exposed to ticks (*Ixodes ricinus* in an active phase: 25 females, 25 nymphs and 10 males) placed in highly air-permeable gauze containers. The containers were fitted onto the forearm and upper arms. Assessments were carried out once every hour for four hours.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



In the cage tests with human volunteers carried out by Todd (1997), 4 µg/cm<sup>2</sup> of formulation containing Saltidin® or DEET were applied onto volunteers forearms (in a 5.65 cm diameter circle). A 20 % Saltidin® formulation provided protection time of 2.5 hours against the deer tick *Ixodes scapularis* (compared to 0.5 hours for DEET)

In the cage tests with human volunteers carried out by Carroll-Loye Biological Research (2010) 0,5g (100 mg Saltidin®) of 20 % formulation were applied onto volunteers arms. A 20 % Saltidin® formulation provided protection time of 14.1 hours against the deer tick *Ixodes scapularis* and 14.0 hours against american dog tick *Dermacentor variabilis* (Table 4).

Table 4: Efficacy (Protection time = PT) of 20 % Saltidin® and DEET formulation against ticks in cage tests.

Test Species	Test Location	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Ixodes ricinus</i>	Near the laboratory Kaiser Wald in Graz, Austria	20.0	4.0**	20.0	4.0	Sixl, 1993 <sup>[9]</sup>
<i>Ixodes scapularis</i>	ICR (Insect Control and Research, Inc.), Baltimore USA	20.0	2.5	20.0	< 0.5	Todd, 1997 <sup>[10]</sup>
<i>Ixodes scapularis</i> (Deer-tick)	USA (California)	20.0	14.1*	Not determined		Carroll-Loye, 2010 <sup>[19]*</sup>
<i>Dermacentor variabilis</i> (American dog tick)	USA (California)	20.0	14.0*	Not determined		Carroll-Loye, 2010 <sup>[19]*</sup>

\* tests carried out on All Family Insect Repellent Spray 20 %

\*\* Duration of the test

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



### 5.3 Effectiveness against Flies

In the efficacy tests carried out against flies, formulations with a low quantity of Saltidin® (7.5 - 12.0) has been used. The repellence against this kind of insect is high even with this amount and even higher with a 20 % Saltidin® content.

Efficacy of formulations containing 7.5 % Saltidin® was determined against tabanids under field conditions. The study was conducted in Austria by Prof. Muelhofer (1993) at the LHS - Labor fuer Hygiene und Sicherheit (Laboratory for Hygiene and Safety). The following abundance of various species was present: *Tabanus bovinus* (horse-fly), 35 %; *Haematopota pluvialis*, 30 %; *Stomoxys calcitrans*, 20 %; and *Chrysops relictus*, 15 %.

Ten volunteers (5/product) wearing protective sleeves with a 10 x 10 cm<sup>2</sup> opening were used in the test. 200 µl of the tested product were applied to each volunteer. The exposure time was 30 minutes and the test was repeated every hour. The test was stopped when one bite occurred over a 30 minute test period. The overall protection time (average of all insect densities) for 7.5 % Saltidin® was five hours and for DEET was six hours (Table 5). At normal insect densities, protection time for 7.5 % Saltidin® was 6 hours and for 10 % DEET was 7 hours. At high insect densities, the corresponding values were 4 and 6 hours. Under extreme insect pressure, the efficacy declined to 4 hours for Saltidin® and to 5 hours for DEET.

Table 5: Efficacy (Protection time = PT) of 7.5 % Saltidin® and 10 % DEET formulations against flies in field tests.

Test Species	Test Location	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Tabanus bovinus</i> , 35 %; <i>Haematopota pluvialis</i> , 30 %; <i>Stomoxys calcitrans</i> , 20 %; <i>Chrysops relictus</i> , 15 %.	Austria	7.5	5.0	10.0	6.0	Muelhofer, 1993 <sup>[11]</sup>

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



In a cage test by Nentwig (1997), a formulation containing 12 % Saltidin® was tested against *Stomoxys calcitrans*. The insects were placed in cages (90 x 30 x 40 cm, with gauze side walls) equipped with two cloth sluices. 90cm<sup>2</sup> of each forearm of a volunteer was treated with 150 µl of the test product. A sleeve with an opening of 25 cm<sup>2</sup> was fastened around the arm: the opening was positioned over the treated area. Both arms were introduced into the cage through the cloth sluice and the number of bites per arm in a three minutes test period was registered.

The protection time for the Saltidin® formulation was 7 hours compared to 4 hours for DEET (Table 6).

Table 6: Efficacy (Protection time = PT) of 12.0 % Saltidin® and 17.0 % DEET formulations against flies in cage tests.

Test Species	Test Location	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Stomoxys calcitrans</i> ,	Bayer AG Lab, Monheim, Germany	12.0	7.0	17.0	4.0	Nentwig, 1997 <sup>[12]</sup>

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## 5.4 Effectiveness against Sandflies

The efficacy of the formulations was tested on the laboratory bred strain of *Phlebotomus (Phlebotomus) duboscqi* on human volunteers (Perrotey *et al.*, 2002, laboratory report and Pesson and Perrotey, 2005, published short communication). A 50 % DEET formulation was also tested for comparative purposes.

In the test, a group of 30 unfed female insects were released into gauze cages (20 x 20 x 20 cm). The test was carried out with six human volunteers (three males and three females). The repellents were applied to the back of the hand and wrist. During the test, a latex glove was worn with a square opening (4 x 4 cm) permitting skin exposure of only a part of hand and wrist. The treated surfaces were exposed in the cage for 5 minutes at hourly intervals until one bite was recorded, terminating the test.

The results of laboratory test with human volunteers showed that Saltidin® formulation had a high degree of repellence against *P. duboscqi* (10 hours). The formulation containing 50 % DEET provided 8.4 hours protection. It can be concluded that Saltidin® is highly effective against Sandflies (Table 7).

Table 7: Efficacy (Protection time = PT) of 20.0 % Saltidin® and 50.0 % DEET formulations against sandflies in cage tests.

Test Species	Test Location	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Phlebotomus (Phlebotomus) duboscqi</i>	Parasitology laboratory, Strasbourg, France	20.0	10	50.0	8.4	Perrotey et al., 2002 <sup>[13]</sup> Pesson and Perrotey, 2005 <sup>[14]</sup>

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## 5.5 Effectiveness against Biting Midges

In a field study with human volunteers, Saltidin® was tested against the Scottish biting midge *Culicoides impunctatus*, and has shown to be a highly effective repellent, preventing landing and biting for more than 8 hours post-application with no significant differences between individuals. It can be concluded that Saltidin® is a highly effective repellent against the Scottish biting midge (*Culicoides impunctatus*).

The test was carried out by Mordue (1999) and Carpenter (2005) at dusk in a marshy field. The application rate was 30 µg of active ingredient/100 cm<sup>2</sup> of skin, the recommended application dose. The average forearm surface area of the 5 volunteers (three males and two females) was 600 cm<sup>2</sup>.

Table 8: Efficacy (Protection time = PT) of 20.0 % Saltidin® formulation against biting midges in field tests.

Test Species	Test Location (Time of test)	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Culicoides impunctatus</i>	Argyllshire, Scotland (dusk)	20.0	> 8.0	Not detected		Mordue, 1999 <sup>[15]</sup> Carpenter, 2005 <sup>[16]</sup>

## 5.6 Effectiveness against Fleas

Saltidin® is recommended as a safe and effective repellent of *Ctenocephalides felis* (cat fleas) for human use.

In the study by Nentwig (1998), 150 µL of 20 % Saltidin® or 20 % DEET products were applied on 90 cm<sup>2</sup> of each arm of two volunteers. Small transparent boxes containing 5 adult fleas each were fastened to the arm treated with the repellent by a bracelet. The number of bites per arm in a three minutes test period was registered, every hour for 9 hours. The repellent effect was considered to fail if two bites occurred in three minutes. It can be concluded that Saltidin® is an effective repellent against fleas.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



Table 9: Efficacy (Protection time = PT) of 20.0 % Saltidin® and DEET formulation against fleas in cage tests.

Test Species	Test Location	Saltidin®		DEET		Study ID/Reference
		(%)	PT (h)	(%)	PT (h)	
<i>Ctenocephalides felis</i>	Bayer Laboratory, Germany	20.0	9.0	20.0	9.0	Nentwig, 1998 <sup>[17]</sup>

## 5.7 Effectiveness against Ants

Saltidin® has excellent barrier properties against ants. The 20% Formulation without perfume was tested by the institute BioGenius GmbH on glazed tiles against ant species *Lasius niger* (Black ant), workers.

The test were carried out with cardboard test boxes (60 x 40 cm) open at the top. A glass bowl is placed in the box with well-moisturised peat and 300 ants in it.

For testing two tiles (15 x 15 cm) are laid in the box on talc-powdered pedestals. One of the tiles is treated with Insect Repellent Spray 20% Saltidin without perfume, the other one is untreated (control). To let the ants get out of the bowl onto the tiles, two bridges made of cardboard are laid between the bowl and the tiles. At the opposite sides of the tiles there are also cardboard bridges which lead to a container with honey. The ant activity is observed by a camera, which is connected to a computer.

The insect repellent spray 20% Saltidin without perfume showed an excellent repellent effect up to the proved test point 7 days after treatment against workers of Black ants.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



Table 10: Repellent efficacy (Protection time = PT) of 20.0 % Saltidin® formulation against ants

Test Species	Test Location	Saltidin®		Study ID/Reference
		(%)	PT (day)	
<i>Lasius niger</i> (Black ant)	BioGenius GmbH Germany	20.0	> 7	Jung, 2009 <sup>[20]</sup>

## 5.8 Effectiveness against Cockroaches

Saltidin® has excellent barrier properties against German and Oriental cockroaches. The 20% Formulation without perfume was tested by the institute BioGenius GmbH on glazed tiles against cockroach species *Blatta orientalis*, 4<sup>th</sup> nymphal stage and species *Blattella germanica*, 4<sup>th</sup> to 5<sup>th</sup> nymphal stage.

The tests were carried out with cardboard test boxes (60 x 40 cm) open at the top. A white cardboard hiding place (12 x 6.5 x 2.5 cm) was placed in it. The cockroaches can pass through a 2 cm wide and 1 cm high opening. Fifty cockroaches were placed in the boxes. An infra-red camera and an infra-red light were placed over the test box. Two glazed tiles were placed in the box. One of the tiles was treated with an amount of product of 10.000mg/m<sup>2</sup>, equivalent to 2.000 mg Saltidin /m<sup>2</sup>. The other tile was untreated. To attract the cockroaches onto the tiles, a crown cap containing porridge oats dissolved in water was placed behind the tiles. The test room was then completely darkened and only the infra-red light left on.

The insect repellent spray 20% Saltidin without perfume showed an excellent/good repellent effect up to the proved test point 7 days after treatment against both species of cockroaches.

Table 11: Repellent efficacy (Protection time = PT) of 20.0 % Saltidin® formulation against cockroaches

Test Species	Test Location	Saltidin®		Study ID/Reference
		(%)	PT (day)	
<i>Blatta orientalis</i> ; <i>Blattella germanica</i>	BioGenius GmbH, Germany	20.0	> 7	Jung, 2009 <sup>[21]</sup>

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



## 6. References

- [1] Yap, H.H. et al. (1997) Progress report of the various tests some of which were published.
- [2] Luepkes, K. H. (2005) Mosquito Repellent Effects of Various Formulations Based on Bayrepel/KBR 3023 against Yellow Fever Mosquito *Aedes aegypti* . Unpublished LANXESS report.
- [3] Zhao, Y. (2008) Repellent effect on mosquitoes. Unpublished LANXESS report.
- [4] Chen, C., Zheng, J., Zhou, J. (2008) Unpublished LANXESS report.
- [5] Yap, H. H., Jahangir, K., Chong, A. C. S., Adanan, C. R., Malik, Y. A., and Rohaizat, B. (1998) Field Efficacy of a New Repellent, KBR 3023, against *Aedes albopictus* (SKUSE) and *Culex quinquefasciatus* in a Tropical Environment. *Journal of Vector Ecology* 23 (1): 62 - 68.
- [6] Frances, S.P., Van dung, N., Beebe, N. W. and Debboun, M. (2002) Field Evaluation of Repellent Formulations against daytime and Nighttime Biting Mosquitoes in a Tropical forest in Northern Australia. *J. Med. Entomology* 39(3): 541 - 544.
- [7] Frances, S.P., Waterson, N.W., Beebe, N.W. and Cooper, R.D. (2004) Field Evaluation of Repellent Formulation Containing DEET and Picaridin against Mosquitoes in Northern Territory, Australia. *J. Med. Entomol.* 41(3); 414 - 417
- [8] Caroll-Loye, (2008) Mosquito Repellent Efficacy Study Report (Protocol LNX-001). Unpublished LANXESS report.
- [9] Sixl, W. (1993) Efficacy study with three preparations called Autan 611, Autan 617 and Autan 619. Geomedizinische Forschungsstelle (Geomedical Research Office); Parasitologische Untersuchungs- und Beratungsstelle (Parasitological Examination and Consulting Agency); Hygiene Institute der Universitat (University Hygiene Institute), Austria. Unpublished LANXESS Report.
- [10] Todd, R. (1997) Evaluation of a personal repellent against deer ticks ICR - Insect Control and Research, Inc. Unpublished LANXESS Report.
- [11] Muehlhofer, (1993) LHS- Labor fuer Hygiene and Sicherheit (Laboratory for Hygiene and Safety, Austria). Unpublished LANXESS report.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.



- [12] Nentwig (1997) Efficacy of a formulation with 12 % KBR 3023 in comparison to a formulation with 17 % Deet on human arms against three different blood sucking diptera. Bayer Report No. 018032.
- [13] Perrotey, S, Madulo-Leblond, G., and Pesson, B. (2002) Laboratory testing of the insect repellent KBR 3023 against *Phlebotomus duboscqi* (Diptera: *Psychodidae*). Parasitol Res (2002) 88:712-713. DOI 10.1007/s00436-002-0635-2
- [14] Pesson, B. and Perrotey, S. (2005) Study of the Repellent Efficacy of BAYREPEL® on *Phlebotomine* Sandflies: testing of new AUTAN® products on laboratory-bred *Phlebotomine* Sandflies. Unpublished LANXESS report.
- [15] Mordue, A.J. (1999) Field evaluation of the repellent KBR 3023 against the Scottish biting midge, *Culicoides impunctatus*. Department of Zoology, University of Aberdeen, Tillydrone Avenue, Aberdeen AB24 2TZ. WHOPS Report.
- [16] Carpenter, S., Eyres, K., McEndrick, L., Smith, L., Mordue, W., and Mordue, A.J. (2005) Parasitology Research, Springer-Verlag 2005, 10.1007/s00436-005-1298-6.
- [17] Nentwig (1998) Efficacy of KBR 3023 in comparison to Deet on human arms against the cat flea, *Ctenocephalides felis*. ID No. 018580 (BG Animal Health)
- [18] Caroll-Loye, (2010) Efficacy of KBR 3023 – with Black Flies under field conditions
- [19] Caroll-Loye, (2010) Efficacy of KBR 3023 – with Ticks under laboratory conditions
- [20] BioGenius GmbH Biology, Germany (2009) Repellent Effect of formulations against Black ants
- [21] BioGenius GmbH Biology, Germany (2009) Repellent Effect of formulations against german cockroaches *Blattella germanica* and oriental cockroaches *Blatta orientalis*
- [22] Alpha-Biocare GmbH Biology, Germany (2010) Efficacy test of Saltidin® formulation and IR3535 formulation against *Culex quinquefasciatus* and *Aedes aegypti*.

This information and our technical advice - whether verbal, in writing or by way of trials - are given in good faith but without warranty, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.